

Virtual Training for Inexperienced Nursing Students in the Operating Room: Reducing Stress while enhancing Knowledge and Self-Efficacy

Inas Redjem ^{a,b}, Arnaud Huault ^b, Jérôme Chapoul ^b, Mathieu Risy ^c, Florian Nouviale ^c, Alexandre Audinot ^c,
Valérie Gouranton ^c, Estelle Michinov ^a, Pierre Jannin ^b

^a LP3C, Université Rennes 2

^b LTSI-Inserm, Université de Rennes

^c INSA Rennes, IRISA, Université de Rennes

The operating room (OR) is a demanding and high-pressure environment, yet, students often have limited exposure during their training, leaving them ill-prepared for clinical practice¹. This study aims to use immersive virtual reality (iVR) to introduce nursing students to the OR environment in a controlled and immersive way, exploring its impact on anxiety, stress, self-efficacy, and knowledge acquisition through an experimental design with pre-post measures.



METHOD

Pedagogical material

We have developed an iVR training tool allowing students to immerse themselves in the OR; navigate through and interact with the environment; obtain detailed information about aseptic areas, medical staff, and OR equipment.

Two conditions

Participants either explored the OR environment using a **slideshow** (n = 16) or through **iVR** (n = 24).




Figure 1. Slideshow (left) and iVR (right) OR experience


Participants

40 second year nursing students **without prior OR experience** were recruited (*M* age = 23; *SD* = 6.5). 87% of participants were female, 32% had previously used VR and 60% were not familiar with video games.

Measures

- **Presence** (Slater-Usuh-Steed Questionnaire)²
- **Anxiety** (STAI 6 items)³
- **Stress** (visual analog scale)
- **Self-efficacy** (9-items visual analog scale)
e.g. 'I feel confident recognizing some of the instruments used by the surgeon during knee surgery'
- **Knowledge** about OR equipment, medical staff and aseptic areas (9-items quiz)

RESULTS




The iVR group reported a significantly **higher level of presence** (immersion), an **increase in anxiety** levels from pre- to post-intervention, and a **reduction in stress** levels post-intervention.

REFERENCES

- ¹ Kayla N. Laraia, et al. 2024. Analysis of Operating Room Personnel Perspectives to Enhance Medical Student Readiness for Surgery. *Journal of Surgical Education* 81, 1, 70-75.
- ² Martin Usuh, Ernest Catena, Arman Sima, and Mel Slater. 2000. Using Presence Questionnaires in Reality. *Presence: Teleoperators and Virtual Environments* 9, 5, 497-503.
- ³ Theresa M. Marteau and Hillary Bekker. 1992. The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *British Journal of Clinical Psychology* 31, 3, 301-306.

This work was supported by a government grant managed by the French National Research Agency under the France 2030 program, reference ANR-21-DMES-0001



The iVR OR experience was **more effective in enhancing self-efficacy**. Both slideshows and iVR equally **enhanced knowledge** without any significant difference between groups.

CONCLUSION

Implementing an iVR initial OR visit offers a practical and immersive solution to address the challenges associated with limited exposure to the OR. By reducing stress while enhancing self-efficacy and knowledge, this innovative approach has the potential to enhance healthcare education and ultimately contribute to improved patient care.